Virginia Surgical Arts Sergio Vendetti MD, DMD

1240 Perimeter Parkway Ste. 401 Virginia Beach, VA 23454

Skin Wound Care Instructions

1.	Do not get present	wound	dressing	wet for an	proximately	24 hours

- 2. After removing dressing, wash gently with 3% hydrogen peroxide (you may use clean gauze pad or Q-tip).
- 3. Apply antibiotic ointment, such Bacitracin or Polysporin, to thinly cover the entire wound area. (Daily application of an antibiotic ointment will minimize risk of infection and help wound heal more evenly with the surrounding skin).
- 4. For facial wounds without stitches: clean twice daily with peroxide and then apply Bacitracin and fresh band-aid for 5-7 days. Do <u>not</u> allow thick scab to form-this will increase scarring.
- 5. Other locations: apply Bacitracin and leave open to air unless the area will be irritated by clothing; in which case apply a band-aid or Telfa dressing. Be sure to remove the dressings before bathing or if it accidentally gets wet.
- 6. Repeat application of 3% hydrogen peroxide and antibiotic ointment <u>twice</u> a day for a total of 5 to 7 days or longer if wound has not sufficiently healed in this time.

1.	If non-dissolva	able statches v	were placed,	they should be remo	ved in:
	() 5 days	() 7 days	() 10 days	() 14 days	

8. After several days, if you notice an increase in tenderness, swelling, redness or pus formation, or any unusual odor, please call our office immediately.

If the wound does not heal to your satisfaction, please call the office and schedule an approintment at (757) 430-7690.

(NOTE: Hydrogen peroxide and Bactracin ointment are available in the pharmacy without a prescription)